

EXTREMAL BEHAVIOUR IN SECTIONAL MATRICES

ANNA BIGATTI, ELISA PALEZZATO, AND MICHELE TORIELLI

ABSTRACT. In this paper we recall the object *sectional matrix* which encodes the Hilbert functions of successive hyperplane sections of a homogeneous ideal. We translate and/or reprove recent results in this language. Moreover, some new results are shown about their maximal growth, in particular a new generalization of Gotzmann's Persistence Theorem, the presence of a GCD for a truncation of the ideal, and applications to saturated ideals.

1. INTRODUCTION

Let K be a field of characteristic 0 and let $P = K[x_1, \dots, x_n]$ be the polynomial ring with n indeterminates and the standard grading. Given a finitely generated \mathbb{Z} -graded P -module $M = \bigoplus_{d \in \mathbb{N}} M_d$, the M_d 's are finite dimensional K -vector spaces. The **Hilbert function** of M , $H_M: \mathbb{Z} \rightarrow \mathbb{N}$ with $H_M(d) := \dim_K(M_d)$, is a very popular and powerful tool of investigation in Commutative Algebra.

From Macaulay [12], it is well known that the computation of the Hilbert function of M may be reduced to the computation of the Hilbert function of some K -algebras of type P/J , where J is a monomial ideal. In particular, if I is a homogeneous ideal in P , then $H_{P/I} = H_{P/\text{LT}(I)}$.

Another very popular practice consists in studying generic hyperplane sections which, in algebraic terms, means quotienting by a generic linear form. The combination of Hilbert functions and hyperplane sections brought to the result by Green [10] (1988).

The *sectional matrix* of an ideal I was introduced by Bigatti and Robbiano in [6] (1997) unifying the concepts of the Hilbert function of a homogeneous ideal I (along the rows) and its hyperplane sections (along the columns). Sectional matrices did not receive much attention, and in this paper we want to revive them. We extend some results in this language, confirming the merit of this tool and suggesting that

Date: April 6, 2017.

2010 Mathematics Subject Classification. Primary 13D40, 05E40, Secondary 13P99 .

Key words and phrases. Sectional Matrix, Hilbert Function, Hyperplane Section, Generic Initial Ideal, Reduction Number, Extremal Behaviour.

further investigation might cast a new light in many aspects of Commutative Algebra.

In Section 2 we set our notation and recall the definition of sectional matrix. In Sections 3 we recall its main properties converting the results from [6] into terms of the quotient P/I (instead of the ideal I). In particular, Theorem 3.11 shows Macaulay's and Green's inequalities. In Section 4, we recall Gotzmann's Persistence Theorem and the sectional matrix analogue from [6]. Moreover, we generalize it into a *sectional* version (Theorem 4.5).

In Section 5, we describe how to deduce information on the dimension and the degree of a homogeneous ideal in terms of the entries of its sectional matrix. In Section 6, we show how the extremal behaviour implies the presence of a GCD for the truncation of homogeneous ideals. In Section 7, we apply these results to the class of saturated ideals. Finally, in Section 8, we present several examples comparing the information given by the sectional matrix, the generic initial ideal, and the resolution of a homogeneous ideals.

The examples in this paper have been computed with CoCoA ([1], [2]).

2. DEFINITIONS AND NOTATION

Let K be a field of characteristic 0 and $P = K[x_1, \dots, x_n]$ be the polynomial ring with n indeterminates with the standard grading. Let $I \subseteq P$ be a homogeneous ideal in P and $A = P/I$. Then A is a graded P -module $\bigoplus_{d \in \mathbb{N}} A_d$, where $A_d = P_d/I_d$.

The definition of the Hilbert function was extended in [6] to the bivariate function encoding the Hilbert functions of the generic hyperplane sections: the **sectional matrix** of a homogeneous ideal I in P . In this paper, we define, in the obvious way, the sectional matrix for the quotient algebra P/I , and then we show how to adapt the results given in [6] to the use of P/I .

Definition 2.1. Given a homogeneous ideal I in $P = K[x_1, \dots, x_n]$, we define the **sectional matrix** of I and of P/I to be the function $\{1, \dots, n\} \times \mathbb{N} \longrightarrow \mathbb{N}$

$$\begin{aligned} \mathcal{M}_I(i, d) &= \dim_K((I + (L_1, \dots, L_{n-i}))/ (L_1, \dots, L_{n-i}))_d, \\ \mathcal{M}_{P/I}(i, d) &= \dim_K(P_d / (I + (L_1, \dots, L_{n-i}))_d), \end{aligned}$$

where L_1, \dots, L_{n-i} are generic linear forms. Notice that $\mathcal{M}_{P/I}(n, d) = H_{P/I}(d)$ and $\mathcal{M}_{P/I}(i, d) = \binom{d+i-1}{i-1} - \mathcal{M}_I(i, d)$.

Remark 2.2. A generic linear form is a polynomial $L = a_1x_1 + \cdots + a_nx_n$ in $K(a_1, \dots, a_n)[x_1, \dots, x_n]$. In this paper we restrict our attention to a field K of characteristic 0, so the equality above holds for any $L' = \alpha_1x_1 + \cdots + \alpha_nx_n$ with $(\alpha_1, \dots, \alpha_n)$ in a non-empty Zariski-open set in \mathbb{P}_K^n . Therefore in this case it is common practice to talk about “generic linear forms in $K[x_1, \dots, x_n]$ ” instead of “ $K(a_1, \dots, a_n)[x_1, \dots, x_n]$ ”

This small example will be used as a running example throughout the paper.

Example 2.3. Let $P = \mathbb{Q}[x, y, z]$ and $I = (x^4 - y^2z^2, xy^2 - yz^2 - z^3)$ an ideal of P . Then the sectional matrix of P/I is

$$\begin{aligned} H_{P/(I+\langle L_1, L_2 \rangle)}(d) = \mathcal{M}_{P/I}(1, d) : & \begin{matrix} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & \dots \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & \dots \end{matrix} \\ H_{P/(I+\langle L_1 \rangle)}(d) = \mathcal{M}_{P/I}(2, d) : & \begin{matrix} 1 & 2 & 3 & 3 & 2 & 1 & 0 & 0 & \dots \end{matrix} \\ H_{P/I}(d) = \mathcal{M}_{P/I}(3, d) : & \begin{matrix} 1 & 3 & 6 & 9 & 11 & 12 & 12 & 12 & \dots \end{matrix} \end{aligned}$$

where the continuations of the lines, which is obvious in this example, will be described in Theorem 4.1.

3. BACKGROUND RESULTS ON SECTIONAL MATRICES

In this section we recall the main properties of sectional matrices from [6] translating them into terms of the quotient P/I . In particular, we describe the persistence theorem and the connection with rgin .

Let σ be a term-ordering on $P = K[x_1, \dots, x_n]$. The **leading term ideal** or **initial ideal** of an ideal $I \subseteq P$ is the ideal generated by $\{\text{LT}_\sigma(f) \mid f \in I \setminus \{0\}\}$. We denote it by $\text{LT}_\sigma(I)$. For any degree-compatible term-ordering σ it is well known that $H_{P/I} = H_{P/\text{LT}_\sigma(I)}$. This nice property does not extend to $\mathcal{M}_{P/I}$, but only one inequality holds, as Conca [7] proved. We write his result in *sectional matrix* notation.

Theorem 3.1. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$ and σ a term-ordering. Then for all $i = 1, \dots, n$ and $d \in \mathbb{N}$ $\mathcal{M}_{P/I}(i, d) \leq \mathcal{M}_{P/\text{LT}_\sigma(I)}(i, d)$.*

Example 3.2. Recall I from Example 2.3. Compare $\mathcal{M}_{P/I}$ with $\mathcal{M}_{P/\text{LT}_\sigma(I)}$, where $\sigma = \text{DegRevLex}$

$$\begin{aligned} \mathcal{M}_{P/\text{LT}_\sigma(I)}(1, d) : & \begin{matrix} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & \dots \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & \dots \end{matrix} \\ \mathcal{M}_{P/\text{LT}_\sigma(I)}(2, d) : & \begin{matrix} 1 & 2 & 3 & 3 & 2 & 1 & 1 & 0 & \dots \end{matrix} \\ \mathcal{M}_{P/\text{LT}_\sigma(I)}(3, d) : & \begin{matrix} 1 & 3 & 6 & 9 & 11 & 12 & 12 & 12 & \dots \end{matrix} \end{aligned}$$

then we observe that $\mathcal{M}_{P/I}(2, 6) = 0 < 1 = \mathcal{M}_{P/\text{LT}_\sigma(I)}(2, 6)$. Notice that the third lines are equal for all degree-compatible term-orderings because the Hilbert functions are the same: $H_{P/I} = H_{P/\text{LT}_\sigma(I)}$.

In this paper we compare some of our results with the ones from [4]. In order to make the comparison clearer to the reader we need to introduce the notion of s -reduction number and to describe how it is stated in terms of the sectional matrix. The definition of s -reduction number has several equivalent formulations and we recall here the one given in [4].

Definition 3.3. Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$. The s -**reduction number**, $r_s(P/I)$, is $\max\{d \mid H_{P/(I+(L_1, \dots, L_s))}(d) \neq 0\}$, where L_1, \dots, L_s are generic linear forms in P . In our language

$$r_s(P/I) = \max\{d \mid \mathcal{M}_{P/I}(n-s, d) \neq 0\}.$$

The **reduction number** $r(P/I)$ is $r_{\dim(P/I)}(P/I)$.

Example 3.4. In Example 2.3 and Example 3.2 we see that I and $\text{LT}_\sigma(I)$ have the same 2-reduction number, $r_2(P/I) = r_2(P/\text{LT}_\sigma(I)) = 2$, and different 1-reduction number: $r_1(P/I) = 5$ and $r_1(P/\text{LT}_\sigma(I)) = 6$. From the equalities $\dim(P/I) = \dim(P/\text{LT}_\sigma(I)) = 1$ it follows that $r(P/I) = 5$ and $r(P/\text{LT}_\sigma(I)) = 6$.

Remark 3.5. Using Theorem 3.1 Conca proved that $r(P/I) \leq r(P/\text{LT}_\sigma(I))$ [7].

Going back to the problem of finding a monomial ideal with the same sectional matrix as P/I , we recall the definitions of *strongly stable* ideal and of *gin*. A monomial ideal J is said to be **strongly stable** if for every power-product $t \in J$ and every i, j such that $i < j$ and $x_j \mid t$, the power-product $x_i \cdot t / x_j$ is in J .

In [8] Galligo proved that, given I a homogeneous ideal in $K[x_1, \dots, x_n]$, with K a field of characteristic 0 and σ a term-ordering such that $x_1 >_\sigma x_2 >_\sigma \dots >_\sigma x_n$, then there exists a Zariski open set $U \subseteq \text{GL}(n)$ and a strongly stable ideal J such that for each $g \in U$, $\text{LT}_\sigma(g(I)) = J$. This ideal is called the **generic initial ideal with respect to σ** of I and it is denoted by $\mathbf{gin}_\sigma(I)$. In particular, when $\sigma = \text{DegRevLex}$, $\mathbf{gin}_\sigma(I)$ is denoted with $\mathbf{rgin}(I)$.

Example 3.6. Consider the ideal $I = (x^4 - y^2z^2, xy^2 - yz^2 - z^3)$ from Example 2.3. Then $\mathbf{rgin}(I) = (x^3, x^2y^2, xy^4, y^6)$.

See [3] for details about the computation of \mathbf{gin} in CoCoA

Remark 3.7. Let I be a homogeneous ideal. If I has a minimal generator of degree d (then also $g(I)$ does), then $\text{rgin}(I)$ has a minimal generator of degree d . The converse is not true in general: consider for example the ideal $I = (z^5, xyz^3)$ in $\mathbb{Q}[x, y, z]$, then $\text{rgin}(I) = (x^5, x^4y, x^3y^3)$ has a minimal generator of degree 6.

The following result represents the sectional matrix analogue of Macaulay's Theorem for Hilbert functions ($H_{P/I} = H_{P/\text{LT}(I)}$): it reduces the study of the sectional matrix of a homogeneous ideal to the combinatorial behaviour of a monomial ideal.

Lemma 3.8. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$. Then $\mathcal{M}_{P/I}(i, d) = \mathcal{M}_{P/\text{rgin}(I)}(i, d) = \dim_K(P_d/(\text{rgin}(I) + (x_{i+1}, \dots, x_n))_d)$.*

Proof. See Lemma 5.5 in [6]. \square

Remark 3.9. Lemma 3.8 shows that when we have a strongly stable ideal J in P (and in particular $\text{rgin}(I)$ is strongly stable) the sectional matrix of P/J is particularly easy to compute because sectioning J by $n-i$ generic linear forms is the same as sectioning J by the smallest $n-i$ indeterminates, x_{i+1}, \dots, x_n .

Using this combinatorial view, [6] proves a combination of Macaulay's and Green's inequalities ([12], [10]) and then the analogue of Gotzmann's Persistence Theorem [9] for sectional matrices.

We recall the definition of binomial expansion following the notation of [6]. If I is a homogeneous ideal then the $(n-1)$ -binomial expansion of $H_I(d)$ corresponds to a "description" of a lex-segment ideal \mathcal{L} in degree d , and the d -binomial expansion of $H_{P/I}(d)$ corresponds to P/\mathcal{L} . See for example [11] Proposition 5.5.13.

Definition 3.10. Let h, i be positive integers. We define the **i -binomial expansion** of h as $h = \binom{h(i)}{i} + \binom{h(i-1)}{i-1} + \dots + \binom{h(j)}{j}$ which is unique requiring that $h(i) > h(i-1) > \dots > h(j) \geq j \geq 1$.

Moreover we define a family of functions related to the expansion in the following way: $(h_i)_t^s := \binom{h(i)+s}{i+t} + \binom{h(i-1)+s}{i-1+t} + \dots + \binom{h(j)+s}{j+t}$.

For short we will write $(h_i)_+^+$ instead of $(h_i)_{+1}^{+1}$ and $(h_i)^-$ instead of $(h_i)^{-1}$.

Here is Theorem 5.6 of [6] and again we convert the statement in terms of the quotient P/I using the properties of the functions derived from the binomial expansion.

Theorem 3.11 (Sectional matrices). *Let I be a homogeneous ideal in the polynomial ring $P = K[x_1, \dots, x_n]$ and $\mathcal{M} := \mathcal{M}_{P/I}$. Then*

- (a) $\mathcal{M}(i, d+1) \leq \sum_{j=1}^i \mathcal{M}(j, d)$ for all $i = 1, \dots, n$ and $d \in \mathbb{N}$.
- (b) (Macaulay) $\mathcal{M}(i, d+1) \leq (\mathcal{M}(i, d)_d)_+^+$ for all $i = 1, \dots, n$ and $d \in \mathbb{N}$.
- (c) $\mathcal{M}(i-1, d) - \mathcal{M}(i-2, d) \leq ((\mathcal{M}(i, d) - \mathcal{M}(i-1, d))_d)_-^-$ for all $i = 3, \dots, n$ and $d \in \mathbb{N}$.
- (d) (Green) $\mathcal{M}(i-1, d) \leq (\mathcal{M}(i, d)_d)_-^-$ for all $i = 2, \dots, n$ and $d \in \mathbb{N}$.

Proof. This is Theorem 5.6 of [6], using the conversions

$$(H_I(d)_{n-1})^+ = (H_{P/I}(d)_d)_+^+ \quad \text{and} \quad (H_I(d)_{n-1})_-^- = (H_{P/I}(d)_d)_-^-$$

(see for example [11] Proposition 5.5.16 and Proposition 5.5.18). \square

Following Macaulay's inequality [5] defined the *maximal growth* for the Hilbert function. We analogously define maximal growth for the sectional matrix following Theorem 3.11.a.

Definition 3.12. Let I be a homogeneous ideal in the polynomial ring P .

- The Hilbert function $H_{P/I}$ has **maximal growth in degree d** if the “Macaulay's identity” holds: $H_{P/I}(d+1) = (H_{P/I}(d)_d)_+^+$.
- The sectional matrix $\mathcal{M}_{P/I}$ has **i -maximal growth in degree d** if the “Bigatti-Robbiano's identity” holds: $\mathcal{M}_{P/I}(i, d+1) = \sum_{j=1}^i \mathcal{M}_{P/I}(j, d)$.

Remark 3.13. For a homogeneous ideal I in $P = K[x_1, \dots, x_n]$ if $\mathcal{M}_{P/I}$ has n -maximal growth in degree d then I , and $\text{rgin}(I)$, have no minimal generators of degree $d+1$.

More precisely, for any $i \in \{1, \dots, n\}$, Corollary 2.7 of [6] implies that $\mathcal{M}_{P/I}$ has i -maximal growth in degree d if and only if $\text{rgin}(I)$ has no minimal generators of degree $d+1$ in x_1, \dots, x_i . (This is a generalization of Lemma 2.17 in [4].)

4. SECTIONAL PERSISTENCE THEOREM

Gotzmann's Persistence Theorem [9] says that, if the generators of an ideal I have degree $\leq \delta$ and the Hilbert function of P/I has maximal growth in degree δ , then it has maximal growth for all higher degrees.

This is also true for sectional matrices. Here we recall the Persistence Theorem 5.8 of [6], and in Theorem 4.5 we will generalize it for i -maximal growth, for $i \leq n$.

Theorem 4.1 (Persistence Theorem [6]).

Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$. If $\mathcal{M}_{P/I}$ has n -maximal growth in degree δ and I has no generators of degree $> \delta$ then it has i -maximal growth for all $i = 1, \dots, n$ and for all degrees $> \delta$.

Moreover, $\mathcal{M}_{P/I}$ has n -maximal growth for all degrees $> \text{reg}(I)$.

Example 4.2. Consider the polynomial ring $P = \mathbb{Q}[x, y, z, t]$ and the ideal $I = (x^4 - x^2yz, x^5 + xy^3z)$ of P . Then

$$\mathcal{M}_{P/I} = \begin{array}{cccccccccc} & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \dots \\ & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ \mathcal{M}_{P/I} = & 1 & 2 & 3 & 4 & 4 & 3 & 2 & 1 & 1 & \dots \\ & 1 & 3 & 6 & 10 & 14 & 17 & 19 & 20 & 21 & \dots \\ & 1 & 4 & 10 & 20 & 34 & 51 & 70 & 90 & 111 & \dots \end{array}$$

Therefore $\mathcal{M}_{P/I}$ has n -maximal growth in degree 7, whereas a direct computation shows that $H_{P/I}$ has n -maximal growth in degree 49.

Remark 4.3. We emphasize that the regularity of a homogeneous ideal I , the highest degree of the generators of $\text{rgin}(I)$, is a much lower number than the highest degree of the generators of the lex-segment ideal with the same Hilbert function of I . This fact makes the following Persistence theorem more “practical” than Gotzmann’s.

In particular, this value is used by CoCoA for truncating the size of the sectional matrix, displaying the rows up to degree $\text{reg}(I) + 1$, so that the last column shows the persisting equalities. In Example 2.3, we have $\text{reg}(I) = 6$ and in degree 7 we read the persisting equalities.

With the next lemma we show that if $\mathcal{M}_{P/I}$ has i -maximal growth in degree δ for some $i < n$, this persists in higher degrees, even if it there is no n -maximal growth.

Lemma 4.4. Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$ generated in degree $\leq \delta + 1$. If there exists $i \leq n$ such that $\text{rgin}(I)$ has no minimal generators of degree $\delta + 1$ in $P_{(i)} = K[x_1, \dots, x_i]$, then $\text{rgin}(I)$ has no minimal generators of any degree $> \delta$ in $P_{(i)}$.

Proof. Let σ be DegRevLex and g a generic change of coordinates. Suppose that the σ -Gröbner basis of $J = g(I)$ has a polynomial f_2 of degree $\delta + 2$. Then f_2 comes from a minimal syzygy of $\text{rgin}(I) = \text{LT}_\sigma(J)$ and hence this syzygy is linear (see Lemma 5.7 in [6]). This means that there exists a minimal generator t_1 of $\text{rgin}(I)$ of degree $\delta + 1$, and, by the hypothesis, all minimal generators of $\text{rgin}(I)$ must be in (x_{i+1}, \dots, x_n) . Let f_1 be the Gröbner basis polynomial such that $t_1 = \text{LT}_\sigma(f_1)$. Because we are using the reverse lexicographic term-ordering, this fact implies that $f_1 \in (x_{i+1}, \dots, x_n)$. As a consequence

any s-polynomial constructed with f_1 is a difference of polynomials in (x_{i+1}, \dots, x_n) , so $f_2 \in (x_{i+1}, \dots, x_n)$. Thus any Gröbner basis element of degree $\delta + 2$ is in (x_{i+1}, \dots, x_n) , and therefore $\text{rgin}(I)$ has no minimal generators of degree $\delta + 2$ in $K[x_1, \dots, x_i]$. Iterating this reasoning, we can conclude that $\text{rgin}(I)$ has no minimal generators of degree $> \delta$ in $K[x_1, \dots, x_i]$. \square

Using Lemma 4.4, we can now extend Theorem 4.1.

Theorem 4.5 (Sectional Persistence Theorem).

Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$ generated in degree $\leq \delta + 1$. If there exists $i \in \{1, \dots, n\}$ such that $\mathcal{M}_{P/I}$ has i -maximal growth in degree δ then it has j -maximal growth for all $j \in \{1, \dots, i\}$ and for all degrees $\geq \delta$.

Proof. From $\mathcal{M}_{P/I}(i, \delta + 1) = \sum_{j=1}^i \mathcal{M}_{P/I}(j, \delta)$ we have that $\text{rgin}(I)$ has no generators of degree $\delta + 1$ in $K[x_1, \dots, x_i]$ (see Remark 3.13). Then, applying Lemma 4.4, we know that $\text{rgin}(I)$ has no generators of degree $> \delta$ in $K[x_1, \dots, x_i]$. Hence we can apply Theorem 4.1 to $J_{(i)} = \text{rgin}(I) \cap K[x_1, \dots, x_i]$ and for all $d > \delta$ and $j = 1, \dots, i$ we get $\mathcal{M}_{P/\text{rgin}(I)}(i, d+1) = \mathcal{M}_{P/J_{(i)}}(j, d+1) = \sum_{k=1}^j \mathcal{M}_{P/J_{(i)}}(k, d) = \sum_{k=1}^j \mathcal{M}_{P/\text{rgin}(I)}(k, d)$. The conclusion now follows from Lemma 3.8. \square

Example 4.6. Consider the polynomial ring $P = \mathbb{Q}[x, y, z, w]$ and the ideal $I = (x^2, xy, xz(z + w), x(z^2 + w^2))$ of P . Then $\text{rgin}(I) = (x^2, xy, xz^2, xzw, xw^3)$. Notice that I is generated in degree ≤ 3 , and its regularity is 4, so $\mathcal{M}_{P/I}$ has n -maximal growth in degree 4. Then

$$\mathcal{M}_{P/I} = \begin{array}{cccccccc} & 0 & 1 & 2 & 3 & 4 & 5 & \dots \\ & 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ \mathcal{M}_{P/I} = & 1 & 2 & 1 & 1 & 1 & 1 & \dots \\ & 1 & 3 & 4 & 4 & 5 & 6 & \dots \\ & 1 & 4 & 8 & 11 & 15 & 21 & \dots \end{array}$$

We see that $\mathcal{M}_{P/I}$ has 2-maximal growth in degree 2 and 3-maximal growth in degree 3:

$$\mathcal{M}_{P/I}(2, 3) = 1 = \mathcal{M}_{P/I}(1, 2) + \mathcal{M}_{P/I}(2, 2)$$

$$\mathcal{M}_{P/I}(3, 4) = 5 = \mathcal{M}_{P/I}(1, 3) + \mathcal{M}_{P/I}(2, 3) + \mathcal{M}_{P/I}(3, 3).$$

Hence from Theorem 4.5 it follows that $\mathcal{M}_{P/I}$ has 3-maximal growth for all degrees ≥ 3 and it has 2-maximal growth for all degrees ≥ 2 .

5. HILBERT POLYNOMIAL, HILBERT SERIES, DIMENSION, MULTIPLICITY

In this section, we show how to read some algebraic invariants for a homogeneous ideal I from its sectional matrix $\mathcal{M}_{P/I}$ truncated at degree δ .

Lemma 5.1. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$ generated in degree $\leq \delta + 1$. If $\mathcal{M}_{P/I}$ has i -maximal growth in degree δ , then for all $d \in \mathbb{N}$ we have*

$$\mathcal{M}_{P/I}(i, \delta+d) = \sum_{j=1}^i \binom{i-j+d-1}{i-j} \cdot \mathcal{M}_{P/I}(j, \delta)$$

Proof. We prove the statement by induction on d . For $d = 1$ the equality holds by hypothesis $\mathcal{M}_{P/I}(i, \delta+1) = \sum_{j=1}^i \mathcal{M}_{P/I}(j, \delta)$.

Now suppose the equality holds for $\mathcal{M}_{P/I}(k, \delta+d)$ for all $k = 1, \dots, i$. Then $\mathcal{M}_{P/I}(i, \delta+d+1) = \sum_{k=1}^i \mathcal{M}_{P/I}(k, \delta+d) = \sum_{k=1}^i \sum_{j=1}^k \binom{k-j+d-1}{k-j} \cdot \mathcal{M}_{P/I}(j, \delta)$ then we swap the sums varying j in $\{1, \dots, i\}$ and $t = k - j$ with $k \in \{j, \dots, i\}$:

$$\sum_{j=1}^i \left(\sum_{t=0}^{i-j} \binom{t+d-1}{t} \right) \cdot \mathcal{M}_{P/I}(j, \delta) = \sum_{j=1}^i \binom{i-j+d-1}{i-j} \cdot \mathcal{M}_{P/I}(j, \delta)$$

and this concludes the proof. \square

Proposition 5.2. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$ generated in degree $\leq \delta + 1$, and let L_1, \dots, L_n be generic linear forms in P . If $\mathcal{M}_{P/I}$ has i -maximal growth in degree δ , then the Hilbert polynomial of $P/(I + (L_1, \dots, L_{n-i}))$ is*

$$p_i(x) = \sum_{j=1}^i \binom{i-j+x-\delta-1}{i-j} \cdot \mathcal{M}_{P/I}(j, \delta)$$

In particular, if $p_i \neq 0$ let $k = \min\{j \in \{1, \dots, i\} \mid \mathcal{M}_{P/I}(j, \delta) \neq 0\}$, then

$$p_i(x) = \frac{\mathcal{M}_{P/I}(k, \delta)}{(i-k)!} x^{i-k} + \dots \text{ lower degree terms.}$$

Proof. The first part of the corollary is trivial from Lemma 5.1: for $x > \delta$ we have $p_i(x) = \mathcal{M}_{P/I}(i, x) = \sum_{j=1}^i \binom{i-j+(x-\delta)-1}{i-j} \cdot \mathcal{M}_{P/I}(j, \delta)$.

We conclude by observing that $\binom{i-k+x-\delta-1}{i-k} = \frac{(x-\delta+i-k-1) \dots (x-\delta+1)(x-\delta)}{(i-k)!}$ and therefore equal to $\frac{1}{(i-k)!} x^{i-k} + (\text{terms of lower degree})$. \square

Proposition 5.3. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$ generated in degree $\leq \delta + 1$, and let L_1, \dots, L_n be generic linear forms*

in P . If $\mathcal{M}_{P/I}$ has i -maximal growth in degree δ , then the Hilbert series of $R_i = P/(I + (L_1, \dots, L_{n-i}))$ is

$$\text{HS}_{R_i}(t) = \sum_{d=0}^{\delta} \mathcal{M}_{P/I}(i, d) t^d + \left(\sum_{j=1}^i \frac{\mathcal{M}_{P/I}(j, \delta)}{(1-t)^{i-j+1}} \right) t^{\delta+1}.$$

Proof. By definition $\text{HS}_{R_i}(t) = \sum_{d=0}^{\infty} \mathcal{M}_{P/I}(i, d) t^d$. By Lemma 5.1, we have that

$$\sum_{d=\delta+1}^{\infty} \mathcal{M}_{P/I}(i, d) t^d = \sum_{d=\delta+1}^{\infty} \left(\sum_{j=1}^i \binom{i-j+d-\delta-1}{i-j} \cdot \mathcal{M}_{P/I}(j, \delta) \right) t^d$$

swapping the sums and letting $k = d - \delta - 1$ it becomes

$$= \sum_{j=1}^i \left(\sum_{k=0}^{\infty} \binom{i-j+k}{i-j} t^k \right) \cdot \mathcal{M}_{P/I}(j, \delta) \cdot t^{\delta+1} = \left(\sum_{j=1}^i \frac{\mathcal{M}_{P/I}(j, \delta)}{(1-t)^{i-j+1}} \right) t^{\delta+1}.$$

Therefore, we can conclude by adding the first part of the series, $\sum_{d=0}^{\delta} \mathcal{M}_{P/I}(i, d) t^d$. \square

The following theorem shows that we can easily read the dimension and the multiplicity of P/I from its sectional matrix. In particular, this information may be found in the δ -th column with $\delta < \text{reg}(I)$ (see Example 5.6).

Theorem 5.4. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$ generated in degree $\leq \delta+1$ such that $I_{\delta} \neq P_{\delta}$ and let $i = \min_j \{\mathcal{M}_{P/I}(j, \delta) \neq 0\}$. If $\mathcal{M}_{P/I}(i, \delta) = \mathcal{M}_{P/I}(i, \delta+1)$, then*

$$\dim(P/I) = n-i+1 \quad \text{and} \quad \deg(P/I) = \mathcal{M}_{P/I}(i, \delta).$$

Proof. The hypothesis implies that $\mathcal{M}_{P/I}$ has i -maximal growth in degree δ , so, by Theorem 4.5, has j -maximal growth in degree d , for all $d > \delta$ and $j = 1, \dots, i$; this means that $i = \min_j \{\mathcal{M}_{P/I}(j, d) \neq 0\}$ for all $d > \delta$ and $\mathcal{M}_{P/I}(i, d) = \mathcal{M}_{P/I}(i, \delta)$ for all $d \geq \delta$. Now, let $\delta' \geq \delta$ such that $\mathcal{M}_{P/I}$ has n -maximal growth in degree δ' , for example $\delta' = \max\{\delta, \text{reg}(I)\}$. Applying Proposition 5.3, and setting the highest power, $(1-t)^{n-i+1}$, as common denominator, it follows that

$$\text{HS}_{P/I}(t) = \frac{\mathcal{M}_{P/I}(i, \delta') + f(t)(1-t)}{(1-t)^{n-i+1}}$$

for some polynomial $f(t) \in K[t]$ and the fraction above is reduced. Therefore, the degree of its denominator, $n-i+1$, is $\dim(P/I)$, and the evaluation of the numerator in 1, $\mathcal{M}_{P/I}(i, \delta)$, is $\deg(P/I)$. \square

Remark 5.5. The hypothesis of Theorem 5.4, are equivalent to the existence of an integer i such that $\mathcal{M}_{P/I}(i, \delta) = \mathcal{M}_{P/I}(i, \delta+1)$ for

$\delta > r_{n-i+1}(P/I)$. This kind of formulation should be more familiar to the readers of [5] and [4].

Example 5.6. Following Example 4.6 we consider $i = 2$, $\delta = 2$ and then $\mathcal{M}_{P/I}(1, 2) = 0$ and $\mathcal{M}_{P/I}(2, 2) = \mathcal{M}_{P/I}(2, 3) = 1 \neq 0$. We then conclude $\dim(P/I) = n-i+1 = 4-2+1 = 3$ and $\deg(P/I) = \mathcal{M}_{P/I}(i, \delta) = \mathcal{M}_{P/I}(2, 2) = 1$.

Note that we deduced this information from the sectional matrix in degree 2, strictly smaller than $\text{reg}(I) = 4$ and also smaller than 3, the maximal degree of the generators of I .

Remark 5.7. For any homogeneous ideal I in $P = K[x_1, \dots, x_n]$, $\mathcal{M}_{P/I}$ has i -maximal growth for all degrees $\geq \text{reg}(I)$ and for all $i \in \{1, \dots, n\}$. Therefore all the results in this section hold replacing their hypotheses with “ $\delta > \text{reg}(I)$ ”.

Example 5.8. In Example 2.3, with $\text{reg}(I) = 6$, we have $i = 3$ so $\dim(P/I) = 1$ and $\deg(P/I) = \mathcal{M}_{P/I}(3, 6) = 12$,

6. MAXIMAL GROWTH AND GREATEST COMMON DIVISOR

Now we make a subtle change: we consider the same scenario in degree δ for an arbitrary homogeneous ideal I and see that the same conclusion hold on the *truncation* $\langle I_{\leq \delta} \rangle$.

Proposition 6.1. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$. If there exists δ such that $I_\delta \neq P_\delta$ and $\mathcal{M}_{P/I}(i, \delta) = \mathcal{M}_{P/I}(i, \delta+1)$, for $i = \min_j \{\mathcal{M}_{P/I}(j, \delta) \neq 0\}$, then we have*

$$\dim(P/\langle I_{\leq \delta} \rangle) = \dim(P/\langle I_{\leq \delta+1} \rangle) = n-i+1,$$

$$\deg(P/\langle I_{\leq \delta} \rangle) = \deg(P/\langle I_{\leq \delta+1} \rangle) = \mathcal{M}_{P/I}(i, \delta).$$

Proof. By construction $\mathcal{M}_{P/I}(j, d) = \mathcal{M}_{P/\langle I_{\leq \delta} \rangle}(j, d)$ for all $j = 1, \dots, n$ and $0 \leq d \leq \delta$ and $\mathcal{M}_{P/I}(j, d) = \mathcal{M}_{P/\langle I_{\leq \delta+1} \rangle}(j, d)$ for all $j = 1, \dots, i$ and $0 \leq d \leq \delta+1$. From Remark 3.13 it follows that $\text{rgin}(I)$ has no minimal generators in degree $\delta+1$ in x_1, \dots, x_i , and therefore nor do $\text{rgin}(\langle I_{\leq \delta} \rangle) \subseteq \text{rgin}(\langle I_{\leq \delta+1} \rangle) \subseteq \text{rgin}(I)$. It then follows that

$$\mathcal{M}_{P/I}(j, \delta+1) = \mathcal{M}_{P/\langle I_{\leq \delta} \rangle}(j, \delta+1) = \mathcal{M}_{P/\langle I_{\leq \delta+1} \rangle}(j, \delta+1) = 0$$

for all $j = 1, \dots, i-1$. This implies that

$$\mathcal{M}_{P/\langle I_{\leq \delta} \rangle}(i, \delta+1) = \mathcal{M}_{P/\langle I_{\leq \delta} \rangle}(i, \delta) \neq 0,$$

$$\mathcal{M}_{P/\langle I_{\leq \delta+1} \rangle}(i, \delta+1) = \mathcal{M}_{P/\langle I_{\leq \delta+1} \rangle}(i, \delta) \neq 0.$$

Now we can apply Theorem 5.4 and get the conclusions. \square

Corollary 6.2. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$. If there exists δ such that $I_\delta \neq P_\delta$ and $\mathcal{M}_{P/I}$ has n -maximal growth in degree δ , let $i = \min\{j > 1 \mid \mathcal{M}_{P/I}(j, \delta) \neq 0\}$, then $\langle I_{\leq \delta} \rangle = \langle I_{\leq \delta+1} \rangle$, $\dim(P/\langle I_{\leq \delta} \rangle) = n-i+1$, and $\deg(P/\langle I_{\leq \delta} \rangle) = \mathcal{M}_{P/I}(i, \delta)$.*

Proof. By hypothesis $\mathcal{M}_{P/I}$ has n -maximal growth in degree δ , hence by Remark 3.13 it follows that I has no minimal generators in degree $\delta+1$, and therefore $\langle I_{\leq \delta} \rangle = \langle I_{\leq \delta+1} \rangle$. Moreover, from Theorem 4.5 it has i -maximal growth in degree δ . Hence we have the equality $\mathcal{M}_{P/I}(i, \delta+1) = \sum_{j=1}^i \mathcal{M}_{P/I}(j, \delta) = \mathcal{M}_{P/I}(i, \delta) \neq 0$ and the conclusion follows from Proposition 6.1. \square

Example 6.3. Consider the polynomial ring $P = \mathbb{Q}[x, y, z, t]$ and the ideal $I = (x^3, x^2y, xy^2, xyz^2, xyz^3)$ of P . Then

$$\mathcal{M}_{P/I} = \begin{array}{cccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & \dots \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 2 & 3 & 1 & 1 & 1 & 1 & 1 & \dots \\ 1 & 3 & 6 & 7 & 7 & 8 & 9 & 10 & \dots \\ 1 & 4 & 10 & 17 & 24 & 32 & 40 & 50 & \dots \end{array}$$

$$\mathcal{M}_{P/I_{\leq 3}} = \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & \dots \\ 1 & 1 & 1 & 0 & 0 & \dots \\ 1 & 2 & 3 & 1 & 1 & \dots \\ 1 & 3 & 6 & 7 & 8 & \dots \\ 1 & 4 & 10 & 17 & 25 & \dots \end{array} \quad \mathcal{M}_{P/I_{\leq 4}} = \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 & \dots \\ 1 & 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 3 & 1 & 1 & 1 & \dots \\ 1 & 3 & 6 & 7 & 7 & 8 & \dots \\ 1 & 4 & 10 & 17 & 24 & 32 & \dots \end{array}$$

We see that $\mathcal{M}_{P/I}$ has $i = 2$ -maximal growth in degree $\delta = 3$. Then by Proposition 6.1, we have that $\dim(P/\langle I_{\leq 3} \rangle) = \dim(P/\langle I_{\leq 4} \rangle) = 3$, and $\deg(P/\langle I_{\leq 3} \rangle) = \deg(P/\langle I_{\leq 4} \rangle) = 1$, regardless what happens in $\mathcal{M}_{P/I}(j, \delta)$ for $j > i = 2$.

The following proposition is the generalization of Proposition 1.6 of [5].

Proposition 6.4. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$. If $\mathcal{M}_{P/I}$ has n -maximal growth in degree δ , then $\text{reg}(\langle I_{\leq \delta} \rangle) \leq \delta$.*

Proof. Let $\bar{I} = \langle I_{\leq \delta} \rangle$. By construction $\mathcal{M}_{P/I}(j, d) = \mathcal{M}_{P/\bar{I}}(j, d)$ for all $j = 1, \dots, n$ and $0 \leq d \leq \delta$. By Remark 3.13, I has no minimal generators in degree $\delta+1$, and hence $\mathcal{M}_{P/I}(j, \delta+1) = \mathcal{M}_{P/\bar{I}}(j, \delta+1)$ for all $j = 1, \dots, n$. This implies that $\mathcal{M}_{P/\bar{I}}$ has n -maximal growth in degree δ , and then, by the Persistence Theorem 4.5, has n -maximal growth in all degrees $> \delta$. By Lemma 3.8, $\mathcal{M}_{P/\bar{I}} = \mathcal{M}_{P/\text{rgin}(\bar{I})}$ and hence by Lemma 4.4, $\text{rgin}(\bar{I})$ has no minimal generators of degree $> \delta$, and then $\text{reg}(\bar{I}) \leq \delta$. \square

In the rest of the section, we generalize some results of [5] and [4] about the existence a common factor when there is a certain kind of maximal growth.

Corollary 6.5. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$. If there exists δ such that $I_\delta \neq \{0\}$ and $\mathcal{M}_{P/I}(2, \delta) = \mathcal{M}_{P/I}(2, \delta + 1)$ (i.e. has 2-maximal growth) then $\langle I_{\leq \delta} \rangle$ has a GCD of degree $\mathcal{M}_{P/I}(2, \delta)$. Furthermore, $\langle I_{\leq \delta+1} \rangle$ shares the same GCD.*

Proof. By Proposition 6.1 $\dim(P/\langle I_{\leq \delta} \rangle) = \dim(P/\langle I_{\leq \delta+1} \rangle) = n-1$ and $\deg(P/\langle I_{\leq \delta} \rangle) = \deg(P/\langle I_{\leq \delta+1} \rangle) = k = \mathcal{M}_{P/I}(2, \delta)$. This means that $\langle I_{\leq \delta} \rangle$ defines a hypersurface of degree k , i.e. $\langle I_{\leq \delta} \rangle = (F) \cap J$ with $\dim(J) < n-1$ and $\deg(F) = k$. Therefore $\langle I_{\leq \delta} \rangle \subseteq (F)$ as claimed. Similarly for $\langle I_{\leq \delta+1} \rangle$. \square

Following the statement of Corollary 6.5, and along the line of ideas in [5], we give a new definition for the potential GCD, based on the sectional matrix instead of the Hilbert function.

Definition 6.6. Let $I \subset P = K[x_1, \dots, x_n]$ be a homogeneous ideal such that $I_\delta \neq \{0\}$. The \mathcal{M} -potential degree of the GCD of I_δ is $k = \mathcal{M}_{P/I}(2, \delta)$.

The following corollary is the generalization of Proposition 2.7 of [5] and of Corollary 5.2 of [4].

Corollary 6.7. *Let I be a homogeneous ideal in $P = K[x_1, \dots, x_n]$ and let δ be such that $I_\delta \neq \{0\}$. Let k be the \mathcal{M} -potential degree of the GCD of I_δ . If $\mathcal{M}_{P/I}$ has i -maximal growth in degree δ for some $i \geq 2$, then $\langle I_{\leq \delta} \rangle$ and $\langle I_{\leq \delta+1} \rangle$ share the same GCD, F , of degree k .*

Proof. By hypothesis $I_\delta \neq \{0\}$, thus $x^d \in \text{rgin}((I_\delta))$ and then we have $\mathcal{M}_{P/I}(2, d) < d+1$. Moreover, $\mathcal{M}_{P/I}$ has i -maximal growth in degree δ , thus in particular has 2-maximal growth in degree δ (by Theorem 4.5). Therefore we conclude by Corollary 6.5. \square

Let us see this result in action on an example.

Example 6.8. Consider the polynomial ring $P = \mathbb{Q}[x, y, z]$ and the ideal $I = (x^3 + y^3, x^2 + 3xy + 2y^2 - xz - yz, x^4 + x^3y, xy^4 - 16xyz^3, y^5 - 3xy^3z - 4y^4z + 12xyz^3 - 25y^3z^2 + 100yz^4)$ of P . Then

$$\mathcal{M}_{P/I} = \begin{array}{cccccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 2 & 2 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 3 & 5 & 6 & 6 & 4 & 3 & 2 & 1 & 1 & \dots \end{array}$$

We see that $\mathcal{M}_{P/I}$ has 2-maximal growth in degree 3 and indeed both I_3 and I_4 have a GCD F of degree $k = \mathcal{M}_{P/I}(2, 3) = \mathcal{M}_{P/I}(2, 4) = 1$. In particular, a direct computation shows that $F = -x - y$.

7. SATURATED IDEALS

A homogeneous ideal I in $P = K[x_1, \dots, x_n]$ is **saturated** if the irrelevant maximal ideal $\mathfrak{m} = (x_1, \dots, x_n)$ is not an associated prime ideal, i.e. $(I : \mathfrak{m}) = I$. For any homogeneous ideal I of P , the **saturation** of I , denoted I^{sat} , is defined by $I^{\text{sat}} := \{f \in P \mid f\mathfrak{m}^\ell \subseteq I \text{ for some integer } \ell\}$.

In this section we apply the results obtained previously to the case of saturated ideals.

Remark 7.1. It is well known that for any homogeneous ideal J there exists $\ell \in \mathbb{N}$ such that $J^{\text{sat}} = J : \mathfrak{m}^\ell$ and therefore $J_d = (J^{\text{sat}})_d$ for all $d \gg 0$.

Remark 7.2. (Bayer-Stillman) Let I be a homogeneous ideal of P . Then $\text{rgin}(I^{\text{sat}}) = \text{rgin}(I)_{x_n \rightarrow 0}$. This shows that if I is saturated, then $\text{rgin}(I)$ has no minimal generators involving x_n .

Lemma 7.3. *Let I be a saturated ideal in $P = K[x_1, \dots, x_n]$. Then $\mathcal{M}_{P/I}$ has n -maximal growth in degree δ if and only if it has $(n-1)$ -maximal growth in degree δ .*

Proof. By Theorem 3.11.(a) n -maximal growth implies $(n-1)$ -maximal growth.

Suppose now $\mathcal{M}_{P/I}$ has $(n-1)$ -maximal growth. By Remark 3.9 this implies that $\text{rgin}(I)$ has no minimal generators in x_1, \dots, x_{n-1} in degree $\delta+1$. Moreover, since $\text{rgin}(I)$ is saturated, from Remark 7.2 there are no minimal generators divisible by x_n . With no minimal generators in degree $\delta+1$ $\mathcal{M}_{P/\text{rgin}(I)}$, and therefore $\mathcal{M}_{P/I}$, has also n -maximal growth. \square

In general the truncation of a saturated ideal is not saturated, as the following example shows. To guarantee that also the truncation is saturated we need some additional hypothesis, see Lemma 7.5.

Example 7.4. Consider the polynomial ring $P = \mathbb{Q}[x, y, z, t]$ and the ideal $I = (yz - xt, z^3 - yt^2, xz^2 - y^2t, y^3 - x^2z, x^3, x^2y^2)$. This ideal is saturated, however, a direct computation shows that the truncation $I_{\leq 3}$ is not saturated.

The following lemma is the generalization of Lemma 1.4 of [5].

Lemma 7.5. *Let I be a saturated ideal in $P = K[x_1, \dots, x_n]$. If $\mathcal{M}_{P/I}$ has $(n-1)$ -maximal growth in degree δ then the ideal $\langle I_{\leq \delta} \rangle$ is saturated.*

Proof. From Lemma 7.3 it follows that $\mathcal{M}_{P/I}$ has n -maximal growth in degree δ .

Let $\bar{I} = \langle I_{\leq \delta} \rangle$ and $\tilde{I} = \bar{I}^{\text{sat}}$. Notice that we have that $\bar{I}_d \subseteq \tilde{I}_d$ for all $d \in \mathbb{N}$. We want to prove that our hypothesis imply $\bar{I}_d = \tilde{I}_d$ for all $d \in \mathbb{N}$.

By Remark 7.1 we have that $\bar{I}_d = \tilde{I}_d$ for all $d \gg 0$.

Let $f \in \tilde{I}_d$ be an element with $d \leq \delta$. Then $f\mathfrak{m}^\ell \subseteq \bar{I}$ for some integer ℓ . Since $\bar{I} \subseteq I$, we have $f\mathfrak{m}^\ell \subseteq I$, and, by hypothesis I is saturated, therefore $f \in I$. Now, I and \bar{I} coincide in degree $\leq \delta$, hence $f \in \bar{I}$. This shows $\bar{I}_d = \tilde{I}_d$ for all $d \leq \delta$.

By Lemma 4.4, I has no minimal generators in degree $\delta+1$, and hence $I_d = \bar{I}_d = \tilde{I}_d$ also for $d = \delta+1$.

By contradiction, let $d > \delta+1$ be the biggest integer such that $\bar{I}_d \subsetneq \tilde{I}_d$. This means that in degree $d+1$

$$\mathcal{M}_{P/\bar{I}}(n, d+1) = \mathcal{M}_{P/\tilde{I}}(n, d+1)$$

and in degree d

$$\mathcal{M}_{P/\bar{I}}(n, d) > \mathcal{M}_{P/\tilde{I}}(n, d)$$

$$\text{and } \mathcal{M}_{P/\bar{I}}(j, d) \geq \mathcal{M}_{P/\tilde{I}}(j, d), \text{ for } j = 1, \dots, n-1.$$

By definition \bar{I} is generated in degree $\delta < d$, hence, by Theorem 4.1, we have that

$$\mathcal{M}_{P/\bar{I}}(n, d+1) = \sum_{j=1}^n \mathcal{M}_{P/\bar{I}}(j, d).$$

Now, using the equalities and inequalities above, we get

$$\mathcal{M}_{P/\bar{I}}(n, d+1) = \mathcal{M}_{P/\tilde{I}}(n, d+1) = \sum_{j=1}^n \mathcal{M}_{P/\tilde{I}}(j, d) > \sum_{j=1}^n \mathcal{M}_{P/\bar{I}}(j, d).$$

This is impossible by the inequalities in Theorem 3.11.(a). \square

Extending Corollary 6.2 to the case of saturated ideals, we can generalize Theorem 3.6 of [4] and Theorem 3.6 of [5].

Corollary 7.6. *Let I be a saturated ideal in $P = K[x_1, \dots, x_n]$. If there exists δ such that $I_\delta \neq P_\delta$ and $\mathcal{M}_{P/I}$ has $(n-1)$ -maximal or n -maximal growth in degree δ , let $i = \min\{j > 1 \mid \mathcal{M}_{P/I}(j, \delta) \neq 0\}$, then $\langle I_{\leq \delta} \rangle$ is a saturated ideal of dimension $n-i$, of degree $\mathcal{M}_{P/I}(i, \delta)$ and it is δ -regular. Moreover, $\dim(P/I) \leq n-i$.*

Proof. By Lemma 7.3, since I is saturated, having $(n-1)$ -maximal or n -maximal growth is equivalent. By Lemma 7.5, $\langle I_{\leq \delta} \rangle$ is a saturated ideal. The conclusions then follows from Corollary 6.2 and Proposition 6.4. \square

Now we can generalize Corollary 5.2 of [4] and Corollary 2.9 of [5].

Corollary 7.7. *Let I be a saturated ideal in $P = K[x_1, \dots, x_n]$. If $\mathcal{M}_{P/I}$ has $(n-1)$ -maximal growth in degree δ and potential degree of the GCD $= k \geq 1$. Then $\langle I_{\leq \delta} \rangle = \langle I_{\leq \delta+1} \rangle$ is saturated and it has a GCD of degree k .*

Proof. From Lemma 7.3 it follows that $\mathcal{M}_{P/I}$ has n -maximal growth in degree δ . Hence Corollary 6.7 and Lemma 7.5 apply. \square

Remark 7.8. Similarly to the previous Corollary 7.6, we might be tempted to extend Proposition 6.1 to the case of saturated ideals, or, equivalently, Corollary 7.6 to the case of 2-maximal growth.

The example below shows that this is not possible.

Example 7.9. As in [4], under the assumption of Proposition 6.1, if I is saturated we can not conclude that $\langle I_{\leq \delta} \rangle$ is saturated. For this example, we consider a first set of 98 points on the conic Q with equation $(z - 3t)(z + 3t) = 0$ in \mathbb{P}^3 and a second set of 16 points outside Q . In this way we obtain a saturated homogeneous ideal I in $P = \mathbb{Q}[x, y, z, t]$ and

$$\mathcal{M}_{P/I} = \begin{array}{cccccccccccccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & \dots \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 2 & 3 & 4 & 5 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 3 & 6 & 10 & 15 & 17 & 13 & 13 & 11 & 9 & 7 & 5 & 3 & 1 & 0 & \dots \\ 1 & 4 & 10 & 20 & 35 & 52 & 65 & 78 & 89 & 98 & 105 & 110 & 113 & 114 & 114 & \dots \end{array}$$

A direct computation shows that $I_{\leq 5}$ is not saturated.

Example 7.10. Consider the polynomial ring $P = \mathbb{Q}[x, y, z, t, h]$ and the strongly stable ideal

$$I = (x^5, x^4y, x^3y^2, x^2y^3, xy^4, x^4z, x^3yz, x^2y^2z, x^4t, xy^3z^2).$$

The ideal I is saturated and

$$\mathcal{M}_{P/I} = \begin{array}{cccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & \dots \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 3 & 4 & 5 & 1 & 1 & 1 & \dots \\ 1 & 3 & 6 & 10 & 15 & 13 & 13 & 14 & \dots \\ 1 & 4 & 10 & 20 & 35 & 47 & 60 & 74 & \dots \\ 1 & 5 & 15 & 35 & 70 & 117 & 177 & 251 & \dots \end{array}$$

We can now apply the previous statements and see that $I_{\leq 5}$ and $I_{\leq 6}$ are saturated and have GCD of degree 1. Moreover, we can check that the GCD is x . For this example [4] does not apply.

8. SECTIONAL MATRICES, GIN, AND RESOLUTIONS

In this section, we will present some examples in order to compare the sectional matrix with other algebraic invariants, such as the Hilbert function H , the generic initial ideal and the minimal resolution.

We can start from two homogeneous ideals with same Hilbert function but different sectional matrix.

Example 8.1. Consider $P = \mathbb{Q}[x, y, z]$ and let

$$I = (x^2, xy, xz, y^3, y^2z, yz^2, z^3) \quad \text{and} \quad J = (x^2, xy, y^2, xz^2, yz^2, z^3)$$

be two ideals in P . Both the ideals are strongly stable and hence, they coincide with their own rgin. These two ideals clearly have distinct rgin, but they have the same Hilbert function (the last row in the sectional matrices). They have different sectional matrix and different graded resolution.

$$\mathcal{M}_{P/I} = \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & \dots \\ 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & 0 & \dots \\ 1 & 3 & 3 & 0 & 0 & \dots \end{array} \quad \mathcal{M}_{P/J} = \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & \dots \\ 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 0 & 0 & 0 & \dots \\ 1 & 3 & 3 & 0 & 0 & \dots \end{array}$$

The resolutions of P/I and P/J are respectively

$$\begin{aligned} 0 \rightarrow P(-4) \oplus P(-5)^3 \rightarrow P(-3)^3 \oplus P(-4)^7 \rightarrow P(-2)^3 \oplus P(-3)^4 \rightarrow P \rightarrow P/I \rightarrow 0. \\ 0 \rightarrow P(-5)^3 \rightarrow P(-3)^2 \oplus P(-4)^6 \rightarrow P(-2)^3 \oplus P(-3)^3 \rightarrow P \rightarrow P/J \rightarrow 0. \end{aligned}$$

In the following example, we show two ideals with the same sectional matrix and same resolution, but different generic initial ideal.

Example 8.2. Consider the polynomial ring $P = \mathbb{Q}[x, y, z]$ and the ideals

$$I = (x^5, x^4y, x^3y^2, x^2y^3, xy^4, x^4z, x^3yz, x^2y^2z, x^3z^2, \underline{x^2yz^2})$$

$$J = (x^5, x^4y, x^3y^2, x^2y^3, xy^4, x^4z, x^3yz, x^2y^2z, x^3z^2, \underline{xy^3z}).$$

Both the ideals are strongly stable and hence, they coincide with their own rgin. These two ideals clearly have distinct rgin, but they have the same Hilbert function, the same sectional matrix and the same resolution.

$$\mathcal{M}_{P/I} = \mathcal{M}_{P/J} = \begin{array}{ccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & \dots \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 & \dots \\ 1 & 2 & 3 & 4 & 5 & 1 & 1 & \dots \\ 1 & 3 & 6 & 10 & 15 & 11 & 12 & \dots \end{array}$$

The resolution of P/I and P/J is

$$0 \rightarrow P(-7)^5 \rightarrow P(-6)^{14} \rightarrow P(-5)^{10} \rightarrow P.$$

In the following example we show two ideals with the same sectional matrix, but different generic initial ideal and different resolution.

Example 8.3. Consider the polynomial ring $P = \mathbb{Q}[x, y, z]$ and the ideals

$$I = (x^5, x^4y, x^3y^2, x^2y^3, xy^4, x^4z, x^2y^2z, x^3z^2, \underline{x^2yz^2})$$

$$J = (x^5, x^4y, x^3y^2, x^2y^3, xy^4, x^4z, x^2y^2z, x^3z^2, \underline{xy^3z}).$$

We have that

$$\text{rgin}(I) = (x^5, x^4y, x^3y^2, x^2y^3, xy^4, x^4z, x^3yz, x^2y^2z, \underline{x^3z^2}, \underline{x^2yz^3}),$$

$$\text{rgin}(J) = (x^5, x^4y, x^3y^2, x^2y^3, xy^4, x^4z, x^3yz, x^2y^2z, \underline{xy^3z}, \underline{x^3z^3}).$$

$$\mathcal{M}_{P/I} = \mathcal{M}_{P/J} = \begin{array}{cccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & \cdots \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & \cdots \\ 1 & 2 & 3 & 4 & 5 & 1 & 1 & 1 & \cdots \\ 1 & 3 & 6 & 10 & 15 & 12 & 12 & 13 & \cdots \end{array}$$

The resolution of P/I is

$$0 \rightarrow P(-7)^2 \oplus P(-8) \rightarrow P(-6)^{11} \rightarrow P(-5)^9 \rightarrow P \rightarrow P/I \rightarrow 0.$$

The resolution of S/J is

$$0 \rightarrow P(-7)^3 \oplus P(-8) \rightarrow P(-6)^{11} \oplus P(-7) \rightarrow P(-5)^9 \rightarrow P \rightarrow P/J \rightarrow 0.$$

In the following example we show two ideals with the same rgin, therefore the same sectional matrix and Hilbert function, but different resolution.

Example 8.4. Consider the polynomial ring $P = \mathbb{Q}[x, y, z]$, and the ideals of P

$$I = (x^4, y^4, z^4, xy^2z^3, x^3yz^2, x^2y^3z) \quad \text{and} \quad J = \text{rgin}(I).$$

These two ideals clearly have the same rgin. However, J has more minimal generators than I , so they have different resolutions.

$$\mathcal{M}_{P/I} = \mathcal{M}_{P/J} = \begin{array}{cccccccccc} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \cdots \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & \cdots \\ 1 & 2 & 3 & 4 & 2 & 0 & 0 & 0 & 0 & \cdots \\ 1 & 3 & 6 & 10 & 12 & 12 & 7 & 0 & 0 & \cdots \end{array}$$

The resolution of P/I is

$$0 \rightarrow P(-9)^7 \rightarrow P(-7)^3 \oplus P(-8)^9 \rightarrow P(-4)^3 \oplus P(-6)^3 \rightarrow P \rightarrow P/I \rightarrow 0.$$

The resolution of P/J is

$$\begin{aligned} 0 \rightarrow P(-8)^5 \oplus P(-9)^7 \rightarrow P(-5)^2 \oplus P(-6)^2 \oplus P(-7)^{10} \oplus P(-8)^{14} \rightarrow \\ \rightarrow P(-4)^3 \oplus P(-5)^2 \oplus P(-6)^5 \oplus P(-7)^7 \rightarrow P \rightarrow P/J \rightarrow 0. \end{aligned}$$

Acknowledgements. The authors thank Professor L. Robbiano for the valuable discussions and for suggesting us Example 7.4.

During the preparation of this paper the third author was supported by the MEXT grant for Tenure Tracking system.

REFERENCES

- [1] J. Abbott, A. M. Bigatti, and G. Lagorio. CoCoA-5: a system for doing Computations in Commutative Algebra. Available at <http://cocoa.dima.unige.it>, 2017.
- [2] J. Abbott and A.M. Bigatti. CoCoALib: a C++ library for doing Computations in Commutative Algebra. Available at <http://cocoa.dima.unige.it/cocoalib>, 2017.
- [3] John Abbott and Anna Maria Bigatti. Gröbner bases for everyone with CoCoA-5 and CoCoALib. arXiv:1611.07306, to appear in the Proc. of the MSJ SI 2015, 2016.
- [4] J. Ahn and J. C. Migliore. Some geometric results arising from the Borel fixed property. *J. Pure Appl. Algebra*, 209(2):337–360, 2007.
- [5] A. M. Bigatti, A. V. Geramita, and J. C. Migliore. Geometric consequences of extremal behavior in a theorem of Macaulay. *Trans. Amer. Math. Soc.*, 346(1):203–235, 1994.
- [6] A. M. Bigatti and L. Robbiano. Borel sets and sectional matrices. *Ann. Comb.*, 1(1):197–213, 1997.
- [7] A. Conca. Reduction numbers and initial ideals. *Proc. Amer. Math. Soc.*, 131(4):1015–1020, 2003.
- [8] A. Galligo. A propos du théoreme de préparation de Weierstrass. In *Fonctions de plusieurs variables complexes*, pages 543–579. Springer, 1974.
- [9] G. Gotzmann. Eine Bedingung für die Flachheit und das Hilbertpolynom eines graduierten Ringes. *Math. Z.*, 158:61–70, 1978.
- [10] M. Green. Restriction of linear series to hyperplanes, and some results of Macaulay and Gotzmann, algebraic curves and projective geometry. In *Proceedings, Trento*, number 1389 in Lecture Notes in Mathematics. Springer, 1988.
- [11] Martin Kreuzer and Lorenzo Robbiano. *Computational commutative algebra 2*. Springer Science & Business Media, 2005.
- [12] F. S. Macaulay. Some properties of enumeration in the theory of modular systems. *Proc. Lond. Math. Soc.*, 26:531–555, 1927.

DEPARTMENT OF MATHEMATICS, GENOA UNIVERSITY, VIA DODECANESO 35,
16146 GENOA, ITALY.

E-mail address: bigatti@dimma.unige.it

DEPARTMENT OF MATHEMATICS, GENOA UNIVERSITY, VIA DODECANESO 35,
16146 GENOA, ITALY.

E-mail address: palezzato@dimma.unige.it

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, KITA 10, NISHI 8,
KITA-KU, SAPPORO 060-0810, JAPAN.

E-mail address: torielli@math.sci.hokudai.ac.jp